




32.)  $U = 1 \text{ kV}$   $d = 5 \text{ cm}$   $\alpha = 60^\circ$   $\vec{B} = ?$   $e^-$   $m = 9,1 \cdot 10^{-31} \text{ kg}$   
 a.)  $\vec{B} \perp \vec{AC}$   $\vec{B} \parallel \vec{AC}$   $q = -e = -1,6 \cdot 10^{-19} \text{ C}$

$W = qU$   $W = \Delta E_k$   $E_k = \frac{1}{2} m v^2$

$\vec{F} = q \vec{v} \times \vec{B}$   $a_{cp} = \frac{v^2}{R}$



$e^-$  sebessége:  $W = eU = \Delta E_k$   
 $eU = \frac{1}{2} m v^2 \Rightarrow \underline{v}$

a.)  $\vec{F} = -e \vec{v} \times \vec{B} \rightarrow \vec{B}$  befelé




$\frac{d/2}{R} = \cos 30^\circ$   
 $R = \frac{d}{2 \cos 30^\circ} = \dots$

$F_c = ma$   
 $e v B = m \frac{v^2}{R}$   
 $\Downarrow$   
 $\underline{B}$



spirál alakú mozgás  
 $T$ : periódus idő  
 $t$ : repülési idő  
 $t = n T_n$  kell legyen  
 sugar  $R_n$



$e v_{\perp} B_n = m \frac{v_{\perp}^2}{R_n}$

$R_n = \frac{m v_{\perp}}{e B_n}$

$\frac{2 R_n \pi}{v_{\perp}} = T_n$

$T_n = 2 \pi \frac{m}{e B_n} \Rightarrow \underline{B_n}$

$v_{\parallel} = v \cos \alpha$   
 $v_{\perp} = v \sin \alpha$

$t = \frac{d}{v_{\parallel}} = \dots$

$T_n = \frac{t}{n} = \dots$